



Fig.2D



Cultured mESC (left) and mNSC (right) by FGF-basic (154 aa), Human (Cat. No. Z03116) combined with LIF, Human (Cat. No. Z02681).

高稳定性


Fig.3



VEGF₁₆₅ Human (Cat. No. Z02689) and IL-3, Mouse (Cat. No. Z03111) showed almost the same ED₅₀ value between different lots.

高品质 (室温放置1个月)

Fig.4




GM-CSF, Human (Cat. No. Z02695) stimulates cell proliferation of TF-1 cells. The ED₅₀ for this effect is less than 0.5 ng/mL.

Activity of GM-CSF, Human (Cat. No. Z02695). ED₅₀=0.11 ng/mL) has no significant change after stored at room temperature for 1 month.

客户发表文章

- Jennifer Brewin, Christoph Mancao, *et al.* Generation of EBV-specific cytotoxic T-cells that are resistant to calcineurin inhibitors for the treatment of post-transplant lymphoproliferative disease. *Blood*. 2009,114(23):4792-803. **IF:9.06** (IL-2, Human, Cat. No. Z03074)
- Sullivan LC1, Clements CS, Beddoe T, *et al.* The heterodimeric assembly of the CD94-NKG2 receptor family and implications for human leukocyte antigen-E recognition. *Immunity*. 2007, 27(6):900-11. **IF:19.795** (Enterokinase, His, Bovine, Cat. No. Z03004)
- Pang SS, Berry R, Chen Z, *et al.* The structural basis for autonomous dimerization of the pre-T-cell antigen receptor. *Nature*. 2010, 467(7317):844-8. **IF:38.597** (Enterokinase, Porcine, Cat. No. Z02199)
- Shimono K, Tung WE, Macolino C, *et al.* Potent inhibition of heterotopic ossification by nuclear retinoic acid receptor- γ agonists. *Nature Medicine*. 2011, 17(4):454-60. **IF:22.864** (BMP-2, Human, Cat. No. Z02913)
- Glen F. Deleavey, Jonathan K. Watts, Tommy Alain, *et al.* Synergistic effects between analogs of DNA and RNA improve the potency of siRNA-mediated gene silencing. *Nucleic acids research*. 2010, 38(13):4547-57. **IF:8.278** (IFN- α 2b, Human, Cat. No. Z03002)
- Sirois M, Robitaille L, Allary R, *et al.* Traf6 And Irf7 Control Hiv Replication In Macrophages. *PLoS One*. 2011, 6(11):e28125. **IF:3.7** (M-CSF, Human, CHO-expressed, Cat. No. Z02924)
- Gao X, Wan F, Mateo K, *et al.* Bacterial effector binding to ribosomal protein s3 subverts NF- κ B function. *PLoS pathogens*. 2009, 5(12):e1000708. **IF:8.136** (TNF- α , Mouse, *P. pastoris*-expressed, Cat. No. Z02918)
- Wang Y, Thomson CA, Allan LL, *et al.* Characterization of pathogenic human monoclonal autoantibodies against GM-CSF. *Proceedings of the National Academy of Sciences of the United States of America*. 2013, 110(19):7832-7. **IF:9.737** (GM-CSF, Human, Cat. No. Z00349)
- Bouchlaka MN, Sckisel GD, Chen M, *et al.* Aging predisposes to acute inflammatory induced pathology after tumor immunotherapy. *The Journal of experimental medicine*. 2013, 210(11):2223-37. **IF:13.214** (GM-CSF, Human, Cat. No. Z00349)
- Moseychuk, O., Akkiraju, H., Dutta, J., *et al.* Inhibition of CK2 binding to BMPRIa induces C2C12 differentiation into osteoblasts and adipocytes. *Journal of cell communication and signaling*. 2013, 7(4):265-278. (BMP-2, Human, Cat. No. Z02913)

请向当地代理商咨询以获得订购信息



重组蛋白产品概览

www.genscript.com.cn

电话: 400-025-8686-5810 025-58897288-5810
地址: 南京市江宁高新园雅熙路28号
PFP150619

邮箱: product@genscript.com.cn
邮编: 211100

金斯瑞生物科技
Make Research Easy

金斯瑞生物科技
Make Research Easy

金斯瑞利用最新的重组表达系统，包括哺乳动物细胞（CHO和HEK 293）、昆虫细胞、酵母和大肠杆菌，进行重组细胞因子、生长因子和趋化因子的生产；目前已开发上市出近400种重组细胞因子产品，并且所有产品都采用无添加剂冻干技术，产品更稳定，彻底避免外源物对实验的干扰。同时，金斯瑞利用强大的哺乳动物细胞表达平台，已经开发了约200种具有完整翻译后修饰、形态结构和活性功能近似天然蛋白的重组细胞因子产品。金斯瑞提供全面的细胞因子，具有极好的批次一致性、卓越的高活性品质和极低的低内毒素水平；金斯瑞的重组细胞因子产品的活性鉴定，全部由国际权威的第三方测活机构出具报告，保证了产品质量的一致性和可靠性。

为科学研究提供品质卓越、具有良好体验的产品，是我们追寻的终极目标！

了解全部产品请查阅：<http://www.genscript.com.cn/products/recombinant-proteins>

产品应用

细胞活性实验
功能性研究
各种细胞的培养
添加物
免疫测定

产品特性

高纯度、高活性
高稳定性、高品质
活性结果由权威第三方机构提供
无菌、无污染
经济优惠、定制化包装

高纯度/高活性/
无添加剂

大肠杆菌/酵母/昆虫/哺乳动物细胞表达系统

经济实惠
定制化大包装

超低的内毒素
($<0.2\text{EU}/\mu\text{g}$)



部分热销产品目录

细胞因子和生长因子

Catalog	Name	Source
Z02913	BMP-2, Human	<i>E. coli</i>
Z00333	EGF, Human	<i>E. coli</i>
Z03066	EGF, Rat	<i>E. coli</i>
Z02975	EPO, Human	CHO
Z03011	FGF-18, Human	<i>E. coli</i>
Z03207	FGF-21, His, Human	CHO
Z03290	FGF-21, Mouse	<i>E. coli</i>
Z02980	G-CSF, Human	CHO
Z03012	GH, Human	<i>E. coli</i>
Z02983	GM-CSF, Human	CHO
Z02979	GM-CSF, Mouse	CHO
Z03003	IFN- α 2a, Human	<i>E. coli</i>
Z02986	IFN- γ , Human	CHO
Z02916	IFN- γ , Mouse	<i>E. coli</i>
Z03017	IGF-I, Human	<i>E. coli</i>
Z03169	IL-1 RA, Human	HEK 293
Z03121	IL-10, Human	CHO
Z03228	IL-17A, His, Human	CHO
Z02922	IL-1 β , Human	<i>E. coli</i>
Z02988	IL-1 β , Mouse	CHO

Catalog	Name	Source
Z02925	IL-4, Human	CHO
Z02996	IL-4, Mouse	CHO
Z02981	IL-5, Human	CHO
Z03262	IL-8 (77aa)/CXCL8, Human	<i>E. coli</i>
Z02971	IP-10/CXCL10, Human	<i>E. coli</i>
Z02924	M-CSF, Human	CHO
Z02930	M-CSF, Mouse	CHO
Z03010	M-CSF, Rat	CHO
Z02529	PDGF-BB, Human	<i>P. pastoris</i>
Z03096	PDGF-BB, Mouse	<i>E. coli</i>
Z02692	SCF, Human	<i>P. pastoris</i>
Z02997	SCF, Mouse	<i>P. pastoris</i>
Z03069	Shh (C24IV), Human	<i>E. coli</i>
Z03050	Shh (C25II), Mouse	<i>E. coli</i>
Z02684	TNF- α , His, Mouse	<i>E. coli</i>
Z01001	TNF- α , Human	<i>E. coli</i>
Z03133	TWEAK, Human	CHO
Z02690	VEGF164, Mouse	<i>P. pastoris</i>
Z02689	VEGF165, Human	<i>P. pastoris</i>
Z03073	VEGF165, Human	HEK 293


工具酶

Catalog	Name	Source
Z03004	Enterokinase, His, Bovine	<i>P. pastoris</i>
Z03376	Enterokinase, His, Bovine (Lyophilized)	<i>P. pastoris</i>
Z01003	Enterokinase, His, Porcine	<i>P. pastoris</i>
Z02199	Enterokinase, His, Porcine (Lyophilized)	<i>P. pastoris</i>

Catalog	Name	Source
Z02799	3C protease, GST	<i>E. coli</i>
Z03092	3C protease, His	<i>E. coli</i>
Z02798	TEV Protease, GST	<i>E. coli</i>
Z03030	TEV Protease, His	<i>E. coli</i>

高纯度


Fig.1




2 μg of BMP-2, Human (Cat. No. Z02913) was resolved with SDS-PAGE under reducing (R) and non-reducing (N) conditions and visualized by Coomassie Blue staining (Left). The purity is $>98\%$ determined by RP-HPLC (Right)

高活性

Fig.2A




BMP-2, Human (Cat. No. Z02913) induce alkaline phosphatase production by C2C12 cells. The ED_{50} for this effect is typically 0.10-0.80 $\mu\text{g}/\text{mL}$.



BMP-2, Human (Cat. No. Z02913) induce alkaline phosphatase production by ATDC5 cells. The ED_{50} for this effect is typically 0.07-0.20 $\mu\text{g}/\text{mL}$.

Fig.2B*



C2C12 cells were treated equally between experiments with 40 pM, 40 nM and 200 nM of BMP2. Mineral deposits and lipid droplet formation were used as defined end points to describe osteoblastic or adipogenic phenotypes, respectively. BMP2 induced osteogenesis independent of concentration(a) and adipogenesis(b).

*数据摘自金斯瑞客户发表的文章^[10]






Fig.2C



FGF-basic (154 aa), Human (Cat. No. Z03116) stimulates cell proliferation of Balb/3T3 cells. The ED_{50} for this effect is 0.1 ng/mL



LIF, Human (Cat. No. Z02681) stimulates cell proliferation of TF-1 cells. The ED_{50} for this effect is 0.09 ng/mL.